



# Exotic quarkonium and spectroscopy




Pavel Krokovny

Budker INP

on behalf of Belle II collaboration

- Introduction
- Observation of  $Y(10753) \rightarrow \omega\chi_{bJ}(1P)$
- $Y(10753) \rightarrow \omega\eta_b(1S)$  and  $\omega\chi_{b0}(1P)$
- $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$
- Summary

# Bottomonium states




Below  $B\bar{B}$  threshold states are well described by the potential models


Above  $B\bar{B}$  threshold states demonstrate unexpected properties:

- hadronic transitions are strongly enhanced (OZI rule violation)
- $\eta$  transitions are not suppressed compare to  $\pi^+\pi^-$  transitions (HQSS violation)
- two charged  $Z_b^+$  states are observed

Hadronic transitions from the states **below the  $B\bar{B}$  threshold** are described by **gluon emission (QCDME)**:



Hadronic transitions from the states **above the  $B\bar{B}$  threshold** can be enhanced due to  **$B\bar{B}$  mesons rescattering**:



$Z_b^+$  states masses coincide with  **$B\bar{B}^*$   $B^*\bar{B}^*$**  thresholds and decays dominantly to constituent mesons:

$Z_b$ decay mode	Branching fraction
$Z_b^+(10610) \rightarrow \Upsilon(nS)/h_b(mP)\pi^+$	$14.4_{-1.9}^{+2.5}\%$
$Z_b^+(10610) \rightarrow B^+\bar{B}^{*0}/\bar{B}^0B^{*+}$	$85.6_{-2.9}^{+2.1}\%$
$Z_b^+(10650) \rightarrow \Upsilon(nS)/h_b(mP)\pi^+$	$26.6_{-4.7}^{+5.0}\%$
$Z_b^+(10650) \rightarrow B^{*+}\bar{B}^{*0}$	$74_{-6}^{+4}\%$

This is a strong indication of the **molecular nature of  $Z_b^+$  states**  
[PRL, 108, 122001 \(2012\)](#)




# Discovery of $Y(10753)$




Observed in the  $e^+e^- \rightarrow Y(nS) \pi^+\pi^-$  cross section energy dependence by Belle [JHEP 10 \(2019\) 220](#)

	$Y(10860)$	$Y(11020)$	New structure
M (MeV/c <sup>2</sup> )	$10885.3 \pm 1.5^{+2.2}_{-0.9}$	$11000.0^{+4.0}_{-4.5} {}^{+1.0}_{-1.3}$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
$\Gamma$ (MeV)	$36.6^{+4.5}_{-3.9} {}^{+0.5}_{-1.1}$	$23.8^{+8.0}_{-6.8} {}^{+0.7}_{-1.8}$	$35.5^{+17.6}_{-11.3} {}^{+3.9}_{-3.3}$



A dip in  $\sigma(e^+e^- \rightarrow b\bar{b})$  can be described by  $Y(10753)$   
[CPC 44, 8, 083001 \(2020\)](#)




24-30 August

Lomonosov 2023



# What is the nature of Y(10753)?



- Far from the thresholds
- Mass does not match Y(3D) theoretical predictions, and D-wave states are not seen in  $e^+e^-$  collisions
- Y(4S) - Y(3D) mixing can be enhanced due to hadron loops

- Conventional  $b\bar{b}$  state:
  - EPJC 80, 1, 59 (2020)
  - PLB 803, 135340 (2020)
  - PRD 101, 1, 014020 (2020)
  - PRD 102, 1, 014036 (2020)
  - PRD 104, 034036 (2021)
  - PRD 105, 074007 (2022)
  - PRD 106, 094013 (2022)
  - EPJP 137, 357 (2022)
- Tetraquark state:
  - CPC 43, 12, 123102 (2019)
  - PLB, 802, 135217 (2020)
- Hadronic molecule with a small admixture of a bottomonium:
  - PRD 103, 074507 (2021)
- Hybrid state:
  - PRD 99, 1, 014017 (2019)






# Study of $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$



JHEP 06 (2021) 137

- $\sigma(e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)})$  has complicated energy dependence spectra, that hard to describe with resonance shapes
- Rescattering and opening of the various  $B\bar{B}$  thresholds cause oscillatory behaviour due to the coupled-channel effect
- Coupled-channel approach is necessary to study  $\sigma(e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)})$  shape




24-30 August

Lomonosov 2023

6

# Global phenomenological analysis



Data:


- Two-body exclusive cross sections  $\sigma(e^+e^- \rightarrow B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)})$
  - Three-body exclusive cross sections  $\sigma(e^+e^- \rightarrow Y(nS) \pi^+\pi^-)$ ,  $n = 1,2,3$
  - $\sigma(e^+e^- \rightarrow h_b(mP) \pi^+\pi^-)$ ,  $m = 1,2$
  - Combined Belle and BaBar  $R_b$  measurement
  - Using coupled-channel approach.
- Poles:  $Y(4S)$ ,  $Y(10753)$ ,  $Y(5S)$  and  $Y(6S)$
- Results: pole positions (mass and width), branching fractions, energy dependence of scattering amplitudes

PRD 106 (2022) 9, 094013



# Requesting more data



- To study  $Y(10753)$  nature
- Improve accuracy below  $Y(5S)$
  
- Perform energy scan at the Belle II experiment
  
- Two Belle II results will be presented:
  - $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$
  - $e^+e^- \rightarrow B\bar{B}, B\bar{B}^*$  and  $B^*\bar{B}^*$





# SuperKEKB collider

- Asymmetric  $e^+e^-$  collider
- Energy limit 11.02 GeV (up to 11.24)
- Luminosity goal:  $6 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$
- Belle II goal: collect  $50 \text{ ab}^{-1}$




$$L = \frac{\gamma_{\pm}}{2er_e} \left( 1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \left( \frac{I_{\pm} \xi_{y\pm}}{\beta_y^*} \right) \left( \frac{R_L}{R_{\xi_{y\pm}}} \right)$$

Higher beam current

Very strong vertical focusing at the interaction point


parameters		KEKB		SuperKEKB		units
		LER	HER	LER	HER	
Beam energy	$E_b$	3.5	8	4	7	GeV
bg		0.425		0.28		
Half crossing angle	$\phi$	11		41.5		mrad
Beta functions at IP	$\beta_x^*/\beta_y^*$	1200/5.9		60/0.3		mm
Beam currents	$I_b$	1.64	1.19	2.5	1.8	A
<b>Luminosity</b>	<b>L</b>	<b><math>2.1 \times 10^{34}</math></b>		<b><math>6.5 \times 10^{35}</math></b>		<b><math>\text{cm}^{-2}\text{s}^{-1}</math></b>

# Belle II detector






# Data taking status



Belle II Online luminosity Exp: 7-26 - All runs





- New luminosity world record  $4.65 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Data taking efficiency is achieved almost 90%.
- Collected more than  $400 \text{ fb}^{-1}$ .

24-30 August

Lomonosov 2023



11

Y(10753) state was observed by Belle in the analysis of the  $e^+e^- \rightarrow Y(nS) \pi^+\pi^-$  ( $n = 1,2,3$ ) cross section energy dependence [JHEP 10 \(2019\) 220](#).



- Belle II collected  $19 \text{ fb}^{-1}$  around Y(10753) in the gaps between Belle energy scan points
- $9.8 \text{ fb}^{-1}$  is taken near Y(10753) peak

# Observation of $Y(10753) \rightarrow \omega\chi_{bJ}(1P)$



- Significant signals are observed in  $\omega\chi_{b1,2}$  final states
- Confirmation of  $Y(10753)$  and observation of its new decay channels
- Order of magnitude difference is observed for this ratio at  $Y(5S)$  and  $Y(10753)$  indicates the difference in the internal structures of these two states:

$$\frac{\sigma(e^+e^- \rightarrow \chi_{bJ}(1P)\omega)}{\sigma(e^+e^- \rightarrow Y(nS)\pi^+\pi^-)} \sim \begin{cases} \sim 1.5 \text{ at } \sqrt{s} = 10.745 \text{ GeV} \\ \sim 0.15 \text{ at } \sqrt{s} = 10.867 \text{ GeV} \end{cases}$$



# Motivation for $Y(10753) \rightarrow \omega \eta_b(1S) / \chi_{b0}(1P)$

- Tetraquark (diquark-antidiquark) interpretation of this state predicts enhancement of  $Y(10753) \rightarrow \eta_b(1S)\omega$  transition: [CPC 43 \(2019\) 12, 123102](#)

$$\frac{\Gamma(\eta_b \omega)}{\Gamma(\Upsilon \pi^+ \pi^-)} \sim 30$$


- Since  $\eta_b(1S)$  does not have easy for reconstruction decay channels, we reconstruct only  $\omega \rightarrow \pi^+ \pi^- \pi^0$  and use its recoil mass to identify the signal:

$$M_{\text{recoil}}(\pi^+ \pi^- \pi^0) = \sqrt{\left(\frac{E_{\text{c.m.}} - E^*}{c^2}\right)^2 - \left(\frac{p^*}{c}\right)^2}$$

- $e^+e^- \rightarrow \omega \chi_{b0}(1P)$  transition was not observed due to  $B[\chi_{b0}(1P) \rightarrow Y(1S)\gamma] = (1.94 \pm 0.27)\%$ ; In charmonium sector  $Y(4220) \rightarrow \chi_{c0}\omega$  decay was found to be enhanced compare to  $Y(4220) \rightarrow \chi_{c1,2}\omega$  by BES III: [PRD 99, 091103\(R\) \(2019\)](#)




# Y(10753) $\rightarrow$ $\omega\eta_b$ (1S) / $\omega\chi_{b0}$ (1P) results



$$\sigma[e^+e^- \rightarrow \omega\chi_{b0}(1P)] < 8.7 \text{ pb}$$

No significant signals are observed, 90% C.L. upper limits are set

This result do not support the prediction within the tetraquark model  
[CPC 43 \(2019\) 12, 123102](#)



$$\sigma[e^+e^- \rightarrow \omega\eta_b(1S)] < 2.5 \text{ pb}$$

$$\text{JHEP 10 (2019) 22: } \sigma[e^+e^- \rightarrow Y(nS)\pi^+\pi^-] \sim 2.0 \text{ pb}$$

# Reconstruction of $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$



$\Delta E'$  vs  $M_{bc}$  at  $E_{cm} = 10.746$  GeV


- Previous Belle analysis: [JHEP 06 \(2021\), 137](#)
- One B meson is fully reconstructed using hadronic channels;
- $B^* \rightarrow B\gamma$  decays are not reconstructed;

$$\Delta E = E_B - E_{cm}/2$$

$$\Delta E' = \Delta E + M_{bc} - m_B$$





- $|\Delta E'| < 18$  MeV; Signal is identified using beam constrained mass:

$$M_{bc} = \sqrt{E_{cm}^2/4 - p_B^2}$$





# $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ signal yields



- Good description of the  $M_{bc}$  in data
- Contribution of  $Y(4S) \rightarrow B\bar{B}$  production via ISR is visible well described by the fit
- E=10.653 GeV sharp cut of the data at right edge  $\Rightarrow$  fast rise of  $B^*\bar{B}^*$  near threshold



# $e^+e^- \rightarrow B^{(*)}\bar{B}^{(*)}$ cross section




Confirming previous Belle result:

- Solid curve – combined Belle + Belle II data fit
- Dashed curve – Belle data fit only

$\sigma(e^+e^- \rightarrow B^*\bar{B}^*)$  rises rapidly above  $B^*\bar{B}^*$  threshold:

- Similar behaviour was seen for  $D^*\bar{D}^*$  cross section PRD 97, 012002 (2018)
- Possible interpretation: resonance or bound state ( $b\bar{b}$  or  $B^*\bar{B}^*$ ) near threshold MPL A 21, 2779 (2006)
- Also explains a narrow dip in  $\sigma(e^+e^- \rightarrow B\bar{B}^*)$  near  $B^*\bar{B}^*$  threshold by destructive interference between  $e^+e^- \rightarrow B\bar{B}^*$  and  $e^+e^- \rightarrow B^*\bar{B}^* \rightarrow B\bar{B}^*$
- $\Upsilon \pi^+\pi^-$  and  $h_b\eta$  final states could also be enhanced PRD 87, 094033 (2013)



# Study of $e^+e^- \rightarrow Y(nS) \pi^+\pi^-$ ( $n = 1, 2, 3$ )






- Previously Belle observed  $Y(10753)$  using combined fit of  $e^+e^- \rightarrow Y(nS) \pi^+\pi^-$  ( $n = 1,2,3$ ) cross section energy dependencies with  $5.2 \sigma$  significance JHEP 10 (2019) 220.
- Belle II:  $Y(10753)$  is observed in the Born cross-section of  $e^+e^- \rightarrow Y(1S,2S) \pi^+\pi^-$ , while no evidence is found in  $e^+e^- \rightarrow Y(3S) \pi^+\pi^-$  channel.

**5.8  $\sigma$**

**10  $\sigma$**

**0.8  $\sigma$**








# Study of $e^+e^- \rightarrow Y(nS) \pi^+\pi^-$ ( $n = 1, 2, 3$ )



- The hint of the  $Y(1S) \pi^+\pi^-$  enhancement at the  $E_{CM} = 10.653$  GeV could correspond to the inelastic decay of a molecular ( $B^*\bar{B}^*$ ) state.
- No signals of intermediate  $Z_b^+(10610)$  or  $Z_b^+(10650)$  resonances are observed.
- Confirmation of  $Y(10750)$





# Conclusion



## Observation of $e^+e^- \rightarrow \omega\chi_{bJ}(1P)$ at $\sqrt{s} = 10.75$ GeV

- $\sigma[e^+e^- \rightarrow \omega\chi_{bJ}(1P)]$  has a peak at 10.75 GeV
- Confirmation of  $Y(10753)$  and observation of its new decay channels

## Search for $e^+e^- \rightarrow \omega\eta_b(1S) / \omega\chi_{b0}(1P)$ at $\sqrt{s} = 10.75$ GeV

- No significant signals are observed
- The upper limit on the  $Y(10753) \rightarrow \eta_b(1S)\omega$  cross-section contradicts the prediction of the tetraquark model

## Energy dependence of $e^+e^- \rightarrow B\bar{B}, B\bar{B}^*$ and $B^*\bar{B}^*$

- Confirmation of “oscillatory” behavior, improvement of the accuracy
- Rapid rise of  $\sigma(e^+e^- \rightarrow B^*\bar{B}^*)$  above threshold - sign for molecular  $B^*\bar{B}^*$  state?

## Study of $e^+e^- \rightarrow Y(1S)\pi^+\pi^-$ ( $n = 1,2,3$ )

- $Y(10753)$  signals are observed in  $Y(1S,2S)\pi^+\pi^-$  channels
- No signals of intermediate  $Z_b$  resonances are observed
- The hint of the  $Y(1S)\pi^+\pi^-$  enhancement at the  $E_{CM}=10.653$

Golden Modes	
$e^+e^- \rightarrow \pi^+\pi^-\Upsilon(pS)(\rightarrow \ell^+\ell^-)$	
$B\bar{B}$ decomposition	Preliminary results
$\pi^+\pi^-$ Dalitz	
$Y_b \rightarrow \omega\eta_b(1S)$	Preliminary result
$Y_b \rightarrow \omega\chi_{bJ}(1P)$	PRL 130, 091902 (2023)
Silver Modes	
$Y_b \rightarrow \pi^+\pi^-X$ (inclusive)	
$Y_b \rightarrow \eta X$ (inclusive)	
$Y_b \rightarrow \eta\Upsilon(1S, 2S)(\rightarrow \ell^+\ell^-)$	
$Y_b \rightarrow \eta'\Upsilon(1S)(\rightarrow \ell^+\ell^-)$	
$Y_b \rightarrow \Upsilon(1S)$ (inclusive)	
Bronze Modes	
$Y_b \rightarrow \gamma X_b$	
$Y_b \rightarrow \pi^0\pi^0\Upsilon(pS)(\rightarrow \ell^+\ell^-)$	
$Y_b \rightarrow KK(\phi)\Upsilon(pS)(\rightarrow \ell^+\ell^-)$	
$Y_b \rightarrow \pi^0\pi^0X$ (inclusive)	
$Y_b \rightarrow \pi^0X$ (incl. or excl.)	
...	