

The Gallium Anomaly

Carlo Giunti

INFN, Torino, Italy



**TWENTY-SECOND LOMONOSOV
CONFERENCE** August, 21-27, 2025
ON ELEMENTARY PARTICLE PHYSICS
MOSCOW STATE UNIVERSITY


10th Anniversary of the 2015 Nobel Prize




- ▶ The Nobel Prize in Physics 2015 was awarded jointly to Takaaki Kajita and Arthur B. McDonald "for the discovery of **neutrino oscillations**, which shows that neutrinos have mass".
- ▶ The **17th International School on Neutrino Physics and Astrophysics** is dedicated to the 10th anniversary of the 2015 Nobel Prize.

Founders of the Theory of Neutrino Oscillations

Bruno Pontecorvo and Samoil Bilenky




Gallium Radioactive Source Experiments



GALLEX

1995 & 1998


$\langle L \rangle_{\text{GALLEX}} \simeq 1.9 \text{ m}$



SAGE

1999 & 2006

$\langle L \rangle_{\text{SAGE}} \simeq 0.6 \text{ m}$




BEST

2021


$\langle L \rangle_{\text{BEST}}^{R1} \simeq 0.7 \text{ m}$

$\langle L \rangle_{\text{BEST}}^{R2} \simeq 1.1 \text{ m}$

Gallium Anomaly (Bahcall Cross Sections)




Before BEST: $\approx 2.3\sigma$ deficit
Mild Anomaly!



After BEST: $\approx 5.9\sigma$ deficit
Strong Anomaly!

SAGE: nucl-ex/0512041, 0901.2200;
Laveder et al: NPPS 168 (2007) 344, hep-ph/0610352,
0711.4222, 1006.3244, 1507.08204;
Kostensalo et al, 1906.10980

BEST: 2109.11482, 2109.14654, 2201.07364;
Barinov and Gorbunov: 2109.14654;
Huber et al: 2111.12530, 2209.02885;
Brdar et al: 2303.05528; Banks et al: 2311.06352;
Schwetz et al: 2303.15524, 306.09422;
Elliott et al: 2303.13623, 2306.03299;
CG-et al: 2209.00916, 2212.09722, 2312.00565, 2507.13103



- ▶ No clear model-independent anomaly from different path lengths.
- ▶ A constant suppression can fit all data.
- ▶ After the BEST measurements, the Gallium Anomaly is still **an anomaly based on the absolute comparison of observed and predicted rates.**


The Ingredients of the Gallium Anomaly

- ▶ The ν_e flux.
- ▶ The ν_e propagation probability.
- ▶ The $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$ detection cross section.
- ▶ The radiochemical ${}^{71}\text{Ge}$ extraction efficiency.

The ν_e Sources



$E \simeq 0.75 \text{ MeV}$ and $E \simeq 0.43 \text{ MeV}$



$E \simeq 0.81 \text{ MeV}$




ν_e Detection

The ν_e Flux

The neutrino flux is evaluated by calorimetric measurements of the activity of the sources:

^{51}Cr Source [SAGE, hep-ph/9803418]




γ energy release per decay:

$$320 \text{ keV} \times 0.099 \simeq 31.7 \text{ keV}$$

Atomic energy release:

$$36.750 \pm 0.084 \text{ keV} \text{ [BEST, 2015]}$$

^{37}Ar Source [SAGE, nucl-ex/0512041]




Atomic energy release:

$$2.751 \pm 0.021 \text{ keV}$$

[SAGE, nucl-ex/0512041]

Atomic Energy Release




Well measured radioactivity of the sources

- ▶ GALLEX Cr1 [PLB 1995] : $1.714^{+0.030}_{-0.043}$ MCi = $63.4^{+1.1}_{-1.6}$ PBq ($\approx 2.5\%$ unc.)
- ▶ GALLEX Cr2 [PLB 1998] : $1.868^{+0.090}_{-0.057}$ MCi = $69.1^{+3.3}_{-2.1}$ PBq ($\approx 3\%$ unc.)
- ▶ SAGE Cr [hep-ph/9803418] : 0.5166 ± 0.0060 MCi = 19.11 ± 0.22 PBq ($\approx 1.2\%$ unc.)
- ▶ SAGE Ar [nucl-ex/0512041] : 0.409 ± 0.002 MCi = 15.1 ± 0.7 PBq ($\approx 4.6\%$ unc.)
- ▶ BEST Cr [arXiv:2109.11482] : 3.414 ± 0.008 MCi = 12.632 ± 0.030 PBq ($\approx 0.3\%$ unc.)

$$(1 \text{ Ci} = 3.7 \times 10^{10} \text{ Bq})$$

The ν_e Propagation Probability



	^{51}Cr				^{37}Ar	
E [keV]	747	752	427	432	811	813
B.R.	0.8163	0.0849	0.0895	0.00930	0.9020	0.098
σ [10^{-46} cm^2]	62.58	63.22	26.72	27.14	71.35	71.63

[Barinov, Cleveland, Gavrin, Gorbunov, Ibragimova, arXiv:1710.06326]


$$\langle P_{\nu_e \rightarrow \nu_e} \rangle = \frac{\int dV L^{-2} \sum_i (\text{B.R.})_i \sigma_i P_{\nu_e \rightarrow \nu_e}(L, E_i)}{\sum_i (\text{B.R.})_i \sigma_i \int dV L^{-2}}$$

[Acero, CG, Laveder, arXiv:0711.4222]

The detection cross section

- Deficit could be due to an **overestimate** of
$$\sigma(\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-)$$

- First calculation: **Bahcall, hep-ph/9710491**




- $\sigma_{\text{G.S.}}$ from $T_{1/2}({}^{71}\text{Ge}) = 11.43 \pm 0.03$ days [Hampel, Remsberg, PRC 31 (1985) 666]

$$\sigma_{\text{G.S.}}({}^{51}\text{Cr}) = (5.54 \pm 0.02) \times 10^{-45} \text{ cm}^2 \quad [\text{Bahcall, hep-ph/9710491}]$$

- $$\sigma({}^{51}\text{Cr}) = \sigma_{\text{G.S.}}({}^{51}\text{Cr}) \left(1 + 0.669 \frac{\text{BGT}_{175}}{\text{BGT}_{\text{G.S.}}} + 0.220 \frac{\text{BGT}_{500}}{\text{BGT}_{\text{G.S.}}} \right)$$

- The contribution of **excited states** is only $\sim 5\%$! [Bahcall, hep-ph/9710491]

The Ground State Cross Section



$$\sigma_{G.S.} \propto \text{BGT}_{G.S.} \propto \Gamma(^{71}\text{Ge}) \propto \frac{1}{T_{1/2}(^{71}\text{Ge})}$$

- ▶ The Gallium Anomaly could be explained with a decrease of the Ground State Cross Section through an increase of the ^{71}Ge half life.
- ▶ Recent measurements of the ^{71}Ge half life: [CG, Li, Ternes, Xin, arXiv:2212.09722]

$$T_{1/2}^{\text{CY}}(^{71}\text{Ge}) = 11.46 \pm 0.04 \text{ d} \quad [\text{Collar and Yoon, 2023}]$$

$$T_{1/2}^{\text{LLNL}}(^{71}\text{Ge}) = 11.468 \pm 0.008 \text{ d} \quad [\text{Norman et al. (LLNL), 2024}]$$

- ▶ Confirm the 1985 Hampel and Remsberg measurement

$$T_{1/2}^{\text{HR}}(^{71}\text{Ge}) = 11.43 \pm 0.03 \text{ d} \quad [\text{Hampel, Remsberg, PRC 31 (1985) 666}]$$

- ▶ Exclude the explanation of the Gallium Anomaly with a decrease of the Ground State Cross Section through an increase of the ^{71}Ge half life.

The $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$ Cross Section

Model	Method	${}^{51}\text{Cr}$			${}^{37}\text{Ar}$		
		$\sigma_{\text{g.s.}}$	σ_{tot}	δ_{exc}	$\sigma_{\text{g.s.}}$	σ_{tot}	δ_{exc}
Bahcall (1997) [hep-ph/9710491]	${}^{71}\text{Ga}(p, n){}^{71}\text{Ge}$	5.53 ± 0.01	$5.81^{+0.21}_{-0.16}$	4.8%	6.62 ± 0.02	$7.00^{+0.49}_{-0.21}$	5.4%
Elliott et al. (2023) [arXiv:2303.13623]	${}^{71}\text{Ga}(p, n){}^{71}\text{Ge} + \text{SM}$	$5.39^{+0.04}_{-0.04}$	$5.69^{+0.28}_{-0.06}$	5.3%	$6.45^{+0.05}_{-0.05}$	$6.85^{+0.35}_{-0.08}$	5.8%
Cadeddu et al. (2025) [arXiv:2507.13103]		$5.41^{+0.07}_{-0.07}$	$5.71^{+0.27}_{-0.08}$	5.3%	$6.47^{+0.08}_{-0.08}$	$6.87^{+0.34}_{-0.10}$	5.8%
Elliott et al. (2023) [arXiv:2303.13623]	${}^{71}\text{Ga}({}^3\text{He}, {}^3\text{H}){}^{71}\text{Ge} + \text{SM}$	$5.39^{+0.04}_{-0.04}$	$5.85^{+0.18}_{-0.10}$	7.9%	$6.45^{+0.05}_{-0.05}$	$7.02^{+0.19}_{-0.14}$	8.1%
Cadeddu et al. (2025) [arXiv:2507.13103]		$5.41^{+0.07}_{-0.07}$	$5.83^{+0.16}_{-0.16}$	7.2%	$6.47^{+0.08}_{-0.08}$	$7.05^{+0.21}_{-0.21}$	8.2%


Units of 10^{-45} cm^2

Size of the Gallium Anomaly

Model	\bar{R}	GA
Bahcall (1997) [hep-ph/9710491]	$0.800^{+0.035}_{-0.038}$	5.9σ
Elliott et al. (2023) [arXiv:2303.13623] (p, n)	$0.817^{+0.028}_{-0.047}$	6.3σ
Cadeddu et al. (2025) [arXiv:2507.13103] (p, n)	$0.814^{+0.029}_{-0.045}$	6.1σ
Elliott et al. (2023) [arXiv:2303.13623] (${}^3\text{He}, {}^3\text{H}$)	$0.795^{+0.030}_{-0.033}$	6.5σ
Cadeddu et al. (2025) [arXiv:2507.13103] (${}^3\text{He}, {}^3\text{H}$)	$0.797^{+0.035}_{-0.034}$	5.3σ

- Large anomaly with all cross section calculations!
- Even neglecting the excited states contribution, the anomaly is larger than 5σ .

Light Sterile Neutrinos



$$L^{\text{osc}} = \frac{4\pi E}{\Delta m^2}$$

Terminology: a eV-scale sterile neutrino
 means: a eV-scale massive neutrino which is mainly sterile

- Minimal perturbation of successful 3ν mixing:
 effective 4ν mixing with $|U_{e4}|, |U_{\mu 4}|, |U_{\tau 4}| \ll 1$

Effective 3+1 Active-Sterile Neutrino Oscillations

Effective short-baseline survival probability of ν_e (Gallium) and $\bar{\nu}_e$ (reactor):

$$P_{ee}^{\text{SBL}} \simeq 1 - \sin^2 2\vartheta_{ee} \sin^2 \left(\frac{\Delta m_{41}^2 L}{4E} \right)$$


with different notations in the literature:

$$\vartheta_{ee} = \vartheta_{14} = \vartheta_{\text{new}} = \vartheta$$

and

$$\Delta m_{41}^2 = \Delta m_{\text{SBL}}^2 = \Delta m_{\text{new}}^2 = \Delta m^2$$

3+1: Gallium Tension with Solar ν and KATRIN



[Cadeddu et al., arXiv:2507.13103]

Solar ν : arXiv:2411.16840; KATRIN: arXiv:2503.18667]

- ▶ Gallium and Solar neutrinos are ν_e .
- ▶ KATRIN and reactor neutrinos are $\bar{\nu}_e$.
- ▶ There is a tension also with reactor antineutrinos.
- ▶ A very exotic CPT violation could remove the tensions with reactor and KATRIN antineutrinos.
[CG, Laveder, arXiv:1008.4750]
- ▶ There is no CPT-violating solution of the Gallium-Solar neutrino tension!

Tentative Alternative Explanations of the Gallium Anomaly

- ▶ Four-Neutrino Mixing (effective $3+1$ Active-Sterile Neutrino Oscillations) is a **general** possibility which can be tested in many experiments and was not invented for the Gallium Anomaly.
- ▶ Typical Alternative Explanations of the Gallium Anomaly are invented **ad hoc** for the Gallium Anomaly.

scenario	comments	our rating
Explanations within the Standard Model		
increased ^{71}Ge half-life (section 2.1 and ref. [39])	would lead to smaller matrix element for $\nu + ^{71}\text{Ga}$; but the ^{71}Ge half-life has been measured many times with different methods in [38], all of which yield consistent results. So it is hard to imagine a bias in these measurements.	★★☆☆☆
new ^{71}Ga excited state (section 2.2)	would imply a bias in the extraction of the $\nu + ^{71}\text{Ga}$ matrix element from the measured ^{71}Ge half-life. Some very old experiments claim the existence of such a state, but this has not been confirmed in more recent observations.	★★☆☆☆
increased $\text{BR}(^{51}\text{Cr} \rightarrow ^{51}\text{V}^*)$ (section 3)	would cause a bias in translating the heat output of the source to a neutrino production rate. Measurements of $\text{BR}(^{51}\text{Cr} \rightarrow ^{51}\text{V}^*)$ show some tension, but it is far less than the shift required to explain the gallium anomaly.	★★★☆☆
^{71}Ge extraction efficiency (section 4)	one of SAGE's calibration runs has revealed a large bias. Could a small, unnoticed, bias have been present in all gallium experiments?	★★★★☆

[Brdar, Gehrlein, Kopp, arXiv:2303.05528]


Explanations beyond the Standard Model

ν_s coupled to ultralight DM (MSW resonance, section 5.1.1)	several exotic ingredients; somewhat tuned MSW resonance; new ν_4 decay channel required for cosmology.	★★★★☆
ν_s coupled to dark energy (MSW resonance, section 5.1.2)	several exotic ingredients; somewhat tuned MSW resonance; cosmology similar to the previous scenario.	★★★☆☆
ν_s coupled to ultralight DM (param. resonance, section 5.1.3)	several exotic ingredients; somewhat tuned parametric resonance; cosmology requires post-BBN DM production via misalignment.	★★★★☆
decaying ν_s (section 5.2)	difficult to reconcile with reactor and solar data; regeneration of active neutrinos in ν_s decays alleviates tension, but does not resolve it.	★★☆☆☆
vanilla eV-scale ν_s (refs. [17, 18])	preferred parameter space is strongly disfavored by solar and reactor data.	★☆☆☆☆
ν_s with CPT violation (ref. [130])	avoids constraints from reactor experiments, but those from solar neutrinos cannot be alleviated.	
extra dimensions (refs. [131–133])	neutrinos oscillate into sterile Kaluza-Klein modes that propagate in extra dimensions; in tension with reactor data.	
stochastic neutrino mixing (ref. [134])	based on a difference between sterile neutrino mixing angles at production and detection (see also [135, 136]); fit worse than for vanilla ν_s .	
decoherence (refs. [137, 138])	non-standard source of decoherence needed; known experimental energy resolutions constrain wave packet length, making an explanation by wave packet separation alone challenging.	
ν_s coupled to ultralight scalar (ref. [139])	ultralight scalar coupling to ν_s and to ordinary matter affects sterile neutrino parameters; can not avoid reactor constraints	

Radiochemical ^{71}Ge Extraction Efficiency

- ▶ BEST outer volume:
10 extractions of ~ 100 ^{71}Ge atoms in $\sim 40\text{ t} \sim 3 \times 10^{29}$ atoms
- ▶ For each experiment, the ^{71}Ge extraction efficiency ϵ^{cal} was obtained with special calibration measurements.
- ▶ An overestimate of the extraction efficiency can obviously explain the Gallium Anomaly:

$$\epsilon^{\text{true}} < \epsilon^{\text{cal}} \implies R^{\text{true}} = \frac{N_{\text{meas}}}{N_{\text{pred}}} \frac{\epsilon_{\text{cal}}}{\epsilon^{\text{true}}} > \frac{N_{\text{meas}}}{N_{\text{pred}}} = R$$



“resolving the anomaly would require the calibration to be off by around 20%”

[Brdar, Gehrlein, Kopp, arXiv:2303.05528]

Checks of the ^{71}Ge Extraction Efficiency

- ▶ GALLEX performed a hot-atom chemistry* test introducing ^{71}As and counting the ^{71}Ge atoms produced in $e^- + ^{71}\text{As} \rightarrow ^{71}\text{Ge} + \nu_e$:
 $100 \pm 1\%$ recovery [GALLEX, PLB 436 (1998) 158]
- ▶ SAGE introduced about $700\ \mu\text{g} \sim 6 \times 10^{18}$ atoms of stable natural Ge carrier at the beginning of each of the 8 exposures:
 $95 \pm 3\%$ recovery [SAGE, arXiv:0901.2200]
- ▶ BEST introduced about $175\ \mu\text{g} \sim 1.5 \times 10^{18}$ atoms of stable natural Ge carrier at the beginning of each of the 10+10 exposures:
 $95 \pm 1.6\%$ recovery [BEST, arXiv:2109.11482]
- ▶ More: see the experimental papers and the review
Elliott, Gavrin, Haxton, arXiv:2306.03299
- ▶ Conclusion: a 20% calibration overestimate seems very unlikely

* The ^{71}Ge recoil energy is close to that in $\nu_e + ^{71}\text{Ga} \rightarrow ^{71}\text{Ge} + e^-$ for a ^{51}Cr source and comparable to the $\sim 3 - 4\ \text{eV}$ chemical binding energy of the extracted GeCl_4 .

Decoherence Explanation of the Gallium Anomaly

[Farzan, Schwetz, arXiv:2306.09422]

- ▶ “Our proposal does not require the presence of sterile neutrinos but implies a modification of the standard quantum mechanical evolution equations for active neutrinos”
- ▶ However, “the decoherence that we postulate here requires exotic new physics”
- ▶ “conventional decoherence based on particle localisation leads only to tiny effects which are negligible for all oscillation experiments considered here”

$$\text{▶ } P_{ee} = \sum_{i=1}^3 |U_{ei}|^4 + \sum_{i \neq j} |U_{ei}|^2 |U_{ej}|^2 e^{-i\phi_{ij}} e^{-\gamma_{ij}L} \quad \text{with} \quad \phi_{ij} = \frac{\Delta m_{ji}^2 L}{2E_\nu} \quad \text{and}$$


$$\gamma_{ij} = \frac{1}{\lambda_{ij}} \left(\frac{0.75 \text{ MeV}}{E_\nu} \right)^r \quad \text{with} \quad \gamma_{23} = \gamma_{12} + \gamma_{13} \pm 2\sqrt{\gamma_{12}\gamma_{13}}$$

- ▶ In Gallium experiments $\phi_{ij} \ll 1 \Rightarrow e^{-i\phi_{ij}} \approx 1$. Best fit: $\lambda_{13} \rightarrow 0 \Rightarrow \gamma_{23} = \gamma_{12}$


$$P_{ee}^{\text{gal}} \approx 1 - \frac{1}{2} \sin^2 2\theta_{13} - \frac{1}{2} \cos^4 \theta_{13} \sin^2 2\theta_{12} (1 - e^{-\gamma_{12}L})$$

- ▶ “Numerically we have $0.5 \cos^4 \theta_{13} \sin^2 2\theta_{12} \approx 0.404$. Hence, we need partial decoherence in the 12 sector to obtain $P_{ee}^{\text{gal}} \simeq 0.8$ ”

	$r = 2$					$r = 12$				
	χ^2_{\min}/dof	$p\text{-val.}$	$\Delta\chi^2$	$\#\sigma$	$\lambda_{12} [\text{m}]$	χ^2_{\min}/dof	$p\text{-val.}$	$\Delta\chi^2$	$\#\sigma$	$\lambda_{12} [\text{m}]$
CS1, BEST	2.0/1	0.16	30.1	5.1	1.44	1.7/1	0.19	30.4	5.2	1.44
CS1, all	7.7/5	0.17	28.6	5.0	1.74	8.3/5	0.14	28.0	4.9	2.10
CS2, BEST	2.6/1	0.11	32.1	5.3	1.19	2.2/1	0.14	32.5	5.4	1.44
CS2, all	8.4/5	0.14	30.0	5.1	1.44	9.2/5	0.10	29.2	5.0	1.74




- ▶ “While this is a huge improvement compared to the p-values of the null hypothesis **the fit is not perfect**”
- ▶ “This is related to the **partial decoherence** in the 12 sector”
- ▶ “It leads to a **distance dependence on the scale of gallium experiments** which in particular predicts different event ratios in the inner and outer detector volumes of the BEST experiment”



► Solar and KamLAND oscillations $\Rightarrow r \gtrsim 10 - 12$ Extreme!

► “A testable prediction of our scenario is a distance dependent deficit at the radioactive source experiments”

Constraints on the decoherence length λ_{21} from reactor rate data using the KI flux model:



- ▶ The shaded region is the approximate preferred region for the explanation of the Gallium Anomaly.
- ▶ At 2σ this explanation of the Gallium Anomaly is not in conflict with reactor rate data for

$$r = -n \gtrsim 7$$

[CG, Ternes, arXiv:2312.00565]

Other Proposed Solutions of the Gallium Anomaly

- ▶ Active-sterile mixing with **wavepacket-induced decoherence**

[Arguelles, Bertolez-Martinez, Salvado, arXiv:2201.05108, Hardin et al, arXiv:2211.02610]

Extremely small wave packet size $\sigma \approx 6.7 \times 10^{-5}$ nm in tension with:

- ▶ a theoretical estimation: $\approx (2 - 14) \times 10^2$ nm [Akhmedov, Smirnov, arXiv:2208.03736]

- ▶ phenomenological bounds: $\sigma > 2.1 \times 10^{-4}$ nm at 90% C.L.

[de Gouvea, De Romeri, Ternes, arXiv:2005.03022, 2104.05806]

In tension with the reactor rates

[CG, Ternes, arXiv:2312.00565]

- ▶ **ν_4 decay**

[Hardin et al, arXiv:2211.02610; Brdar, Gehrlein, Kopp, arXiv:2303.05528]

In tension with the reactor rates

[CG, Ternes, arXiv:2312.00565]

- ▶ **Broad ν_4 mass distribution**

[Banks, Kelly, McCullough, Zhou, arXiv:2311.06352]

In tension with the reactor rates

[CG, Ternes, arXiv:2312.00565]

Compatibility with Solar Neutrinos

- ▶ The Gallium experiments SAGE and GALLEX/GNO measured the solar neutrino flux with about 5% uncertainty (66.1 ± 3.1 SNU)
- ▶ Taking into account the results of other solar neutrino experiments and neutrino oscillations

[SAGE, arXiv:0901.2200]

$$\Phi_{pp}^{\text{Gallium}} = (6.0 \pm 0.8) \times 10^{10} \text{ cm}^{-2} \text{ s}^{-1}: \text{ about } 13\% \text{ uncertainty}$$

- ▶ In agreement with the Borexino measurement

[Borexino, Nature 562 (2018) 505]

$$\Phi_{pp}^{\text{Borexino}} = (6.1 \pm 0.5^{+0.3}_{-0.5}) \times 10^{10} \text{ cm}^{-2} \text{ s}^{-1}: \text{ about } 12\% \text{ uncertainty}$$

and with the Standard Solar Model (SSM) predictions

$$\Phi_{pp}^{\text{HZ}} = 5.98(1 \pm 0.006) \times 10^{10} \text{ cm}^{-2} \text{ s}^{-1}$$


$$\Phi_{pp}^{\text{LZ}} = 6.03(1 \pm 0.005) \times 10^{10} \text{ cm}^{-2} \text{ s}^{-1}$$

[Vinyoles et al, arXiv:1611.09867]

- ▶ The $\sim 20\%$ reduction of the detection efficiency of the Gallium experiments indicated by the Gallium Anomaly, which implies a $\sim 20\%$ increase of $\Phi_{pp}^{\text{Gallium}}$, would be in tension with Borexino and SSM

Global Fit of Solar Neutrino Data

[Gonzalez-Garcia, Maltoni, Pinheiro, Serenelli, arXiv:2311.16226]




- ▶ f_{GA} multiplies the predicted event rates of all solar fluxes in the Gallium experiments
- ▶ “the global analysis of the solar experiments **do not support** a modification of the neutrino capture cross section in Gallium (or any other effect inducing an energy-independent reduction of the detection efficiency in the Gallium experiments)”
- ▶ “This is so because the global fit implies a rate of pp and ${}^7\text{Be}$ neutrinos in the Gallium experiment which is **in good agreement** with the luminosity constraint as well as with the rates observed in Borexino”

Future Proposals

- ▶ Perform another high-intensity experiment with ^{51}Cr or another source
- ▶ BEST-2: 50 tons of ^{71}Ga divided into 3 zones, irradiated with a ^{65}Zn EC source

[Gorbachev, Gavrin, Ibragimova, Phys.Atom.Nucl. 86 (2023) 1385]



Sensitivity for a 0.5 MCi ^{65}Zn source

- ▶ Longer half life: 244.01 ± 0.09 d
Allows more extractions
- ▶ Higher ν_e energy: 1350 keV
Probes larger values of Δm_{41}^2
- ▶ Larger detection cross section
Bigger event rate
- ▶ **Problem:** larger contribution of model-dependent transitions to higher energy excited states of ^{71}Ge

- ▶ Real-time detection of ν_e from a ^{51}Cr source with a Cerium-doped Gadolinium Aluminum Gallium Garnet (Ce:GAGG) **crystal scintillator detector**: [Huber, arXiv:2209.02885]
 - ▶ $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$ and $\nu_e + e^- \rightarrow \nu_e + e^-$
 - ▶ “With 1.5 tons of scintillator and 10 **source runs** of 3.4 MCi, each, we obtain about 1700 gallium capture events with a purity of 90% and 680,000 neutrino electron scattering events”
 - ▶ “this configuration would allow us to test the gallium anomaly at more than 5σ in an independent way”

- ▶ A strong ν_e or $\bar{\nu}_e$ source inside or near a **liquid scintillator detector** as in the cancelled projects SOX with the Borexino detector [SOX, arXiv:1304.7721] and CeLAND with the KamLAND detector [arXiv:1312.0896] :
 - ▶ **JUNO**: [arXiv:1507.05613]
 - ▶ A monochromatic ^{51}Cr or ^{37}Ar source of ν_e detected with $\nu_e + e^- \rightarrow \nu_e + e^-$
 - ▶ A ^{144}Ce - ^{144}Pr source of $\bar{\nu}_e$ with a continuous β spectrum detected with the Inverse Beta Decay (IBD) reaction $\bar{\nu}_e + p \rightarrow n + e^+$
 - ▶ A cyclotron-produced ^8Li source (**IsoDAR**) of $\bar{\nu}_e$ with a continuous β spectrum detected with the IBD reaction
 - ▶ A ^{51}Cr source in JUNO or JNE [Ciuffoli, Evslin, Gao, Lin, Tang, arXiv:2504.16590]

- ▶ Real-time detection of ν_e from a ^{51}Cr source with a ^{115}In target (as in the old LENS project) dissolved in a **liquid scintillator detector** (as LiquidO):
[Chauhan and Huber, arXiv:2507.07397]
 - ▶ $\nu_e + ^{115}\text{In} \rightarrow ^{115}\text{Sn}^* + e^-$ and $^{115}\text{Sn}^* \rightarrow ^{115}\text{Sn} + 2\gamma$
 - ▶ “a 100 ton indium target combined with 2 source runs of a 3.4 MCi ^{51}Cr source can probe the complete parameter space of the gallium anomaly”

Conclusions

- ▶ Light Sterile Neutrinos can be powerful messengers of BSM New Physics.
- ▶ Historically, the existence of light sterile neutrinos is motivated by the LSND, Gallium, and Reactor Short-Baseline Anomalies.
- ▶ The Reactor Antineutrino Anomaly, discovered in 2011, is fading away.
- ▶ The Gallium Neutrino Anomaly, discovered in 2007, has been revived by the BEST results. It is now the most significant anomaly in neutrino physics.
- ▶ As in 2010, before the discovery of the Reactor Antineutrino Anomaly in 2011, there is a Reactor Antineutrino–Gallium Neutrino tension.
- ▶ There is also a tension with solar neutrinos and KATRIN antineutrinos.
- ▶ No convincing SM explanation of the Gallium Anomaly has been found.
- ▶ Difficulty: known BSM explanations of the Gallium Anomaly affect also solar and reactor neutrinos if not fine-tuned with ad-hoc assumptions.