



# *ROTATION CURVES OF OUR GALAXY AND NEW PARTICLES AS CANDIDATES FOR THE ROLE OF DARK MATTER PARTICLES*

**A.T. D'yachenko<sup>1,2</sup>**

<sup>1</sup>*Emperor Alexander I Petersburg State Transport University;*

<sup>2</sup>*NRC "Kurchatov Institute" B.P. Konstantinov Petersburg Nuclear  
Physics Institute*

*. The search for new particles beyond the Standard Model at accelerators is undoubtedly one of the main tasks of modern nuclear physics. On the other hand, the interpretation of the spectra of soft photons in collisions of high-energy elementary particles is also a mystery for elementary particle physics.*

*Using the black body formula, it is possible to describe both the spectra of strange particles and the spectra of new particles X17 and X38 with the found temperature already for a “cold” electromagnetic plasma, when compared with experimental data for soft photons.*

*As is known, rotation curves of rotation of stars of our Galaxy - the Milky Way and other galaxies cannot be explained without involving hidden mass - dark matter. This is currently one of the main issues of modern Natural Science. A possible interpretation of the increasing speed with distance from the center of the Galaxy is the inclusion of a dark matter halo*

*In this paper, within the framework of the equations of non-relativistic hydrodynamics for dark matter, taking into account the gravitational interaction, we obtained an approximate solution to the problem.*

II

**... важно сконцентрировать имеющиеся ресурсы на основных прорывных направлениях..." В.В. Путин**



*In [1-3], we have shown the manifestation of new  $X17$  and  $X38$  particles with masses of 17 MeV and 38 MeV from the spectra of soft photons in collisions of protons and nuclei at high energies. They were discovered earlier in the experiments of the Hungarian ATOMKI group (boson  $X17$ ) and in collisions of protons and high-energy nuclei at the International Nuclear Centers CERN (Switzerland) (boson  $X17$ ) and JINR (Dubna) (boson  $X38$ ). These new particles can be considered dark matter particles. Recall that the visible part of the mass of the Universe is only 5% in the Universe. The obtained indications of the existence of these new particles need independent confirmation.*

- ✓ *That is, such an interpretation of the spectrum of soft photons (its hardening) can serve as another evidence in favor of the existence of a new particle, the  $X17$  boson. On the same Figure 1 also shows the contribution of photons from the decay of another new particle, the  $X38$  boson with a mass of 38 MeV (dash-dotted line), predicted in experiments carried out in Dubna [5]. However, its contribution appears only at the tail of the spectrum and is weakly visible in the figure. Note that such a hardening of the photon spectrum is not obtained due to the contribution of photons from the hot spot. Since in this case its contribution is 10 times less than the total cross section, and the photon mass is zero, the hardening of the photon spectrum in the energy region of 20 MeV due to the higher hot spot temperature is still negligibly small.*



# LHC

$$\frac{dN}{dp_T} = CTp_T \exp\left(-\frac{\sqrt{m^2 + p_T^2} - m}{T}\right) \quad T = \left(\frac{E_0}{g_Q V_R} 10^9\right)^{1/4} \quad g_Q = (2 \times 8 + \frac{7}{8} 2 \times 2 \times 3 \times 3) = 47$$




# Dark matter for soft photons

$$p + p \rightarrow 2\gamma + X$$

$$M^2 = 2\pi\rho n$$

$$p + C \rightarrow 2\gamma + X$$



On Figure 1 shows the experimental spectrum of soft photons (points [4]), as well as the calculation with allowance for the X17 boson (solid line) and without it (dashed line), the dash-dotted line is the contribution from the X17 boson. It can be seen from the figure that, without taking into account the contribution from the X17 boson, the calculation underestimates the experimental data, and taking this contribution into account reproduces them

A. Belogianni et al., Phys. Lett. B548, 129 (2002)

K. Abraamyan et al., EPJ Web of Conf. 204, 08004 (2019)




A.J. Krasznahorkay et al., Phys. Rev. Lett. 116, 042501 (2016)

C.Y. Wong, JHEP, 08165 (2020); Eur. Phys. J. A, 58 (2022) 100

A.T. D'yachenko, Phys. Atom. Nucl. 83, 1597 (2020); Phys. Atom. Nucl. 85, 1028(2022)



## *X17 and E38*



$$T^3 V = T_0^3 V_0$$

$$\frac{dN}{dp_T} = C p_T \sqrt{p_T} \exp\left(-\frac{p_T}{T}\right)$$

$$\frac{dN}{dp_T} = C T \sqrt{p_T} \left( (E_- + T) \exp\left(\frac{\mu - E_-}{T}\right) - (E_+ + T) \exp\left(\frac{\mu - E_+}{T}\right) \right)$$

$$E_{\pm} = \pm p_1 + m$$

## ***Rotation Curves***

- In the hydrodynamic continuity equation, we included a term of dark matter annihilation proportional to  $A\rho^2$ , where  $\rho$  is the particle density, the particle annihilation coefficient is equal to

$$A = \langle \sigma v \rangle / m_D, \quad (1)$$

- where  $\langle \sigma v \rangle$  is the annihilation cross-section,  $m_D$  is the mass of dark matter particles [8,9]. As a result, we consider a system of hydrodynamic equations for dark matter in spherical coordinates, where equation (2) is the continuity equation for the density  $\rho$ , taking into account the annihilation of dark matter, equation (3) is the Euler equation for the velocity  $\vec{v}$ , taking into account the gravitational potential  $\varphi$ , for which we use the Poisson equation, where  $G$  is the gravitational constant

$$\frac{\partial \rho}{\partial t} + \frac{\partial(r^2 \rho v)}{r^2 \partial r} = -A\rho^2, \quad (2) \quad \frac{\partial \vec{v}}{\partial t} + \vec{v} \nabla \vec{v} = -\nabla \varphi, \quad (3)$$

$$\frac{1}{r^2} \frac{d}{dr} \left( r^2 \frac{d\varphi}{dr} \right) = 4\pi G \rho, \quad (4)$$



## Equations

The velocity  $\vec{v} = \vec{v}_r + \vec{v}_\tau$  can be represented as radial and tangential velocities, respectively. Assuming that the distribution function of dark matter particles is isotropic, we can assume that  $v_r = v_\tau = v$ . In addition,  $\rho(r,t) = \rho(r)$ ,  $v(r,t) = v(r)$ , and  $\varphi(r,t) = \varphi(r)$  are independent of time. From equation (3) we find  $\frac{d(v_\tau \vec{e}_\tau)}{dt} = v_\tau \frac{d\vec{e}_\tau}{dt} = -\frac{v_\tau^2}{r} \vec{e}_r = -\frac{d\varphi}{dr} \vec{e}_r$  Therefore

$$\frac{v_\tau^2}{r} = \frac{d\varphi}{dr} \quad (5) \quad \frac{d(rv^2)}{dr} = 4\pi G\rho, (6) \quad \frac{1}{r^2} \frac{d(r^2 \rho v)}{\partial r} = -A\rho^2. (7)$$

$$\frac{d}{dr} \left( v \left( \frac{d(rv^2)}{dr} \right) \right) = -\frac{A}{4\pi G r^2} \left( \frac{d(rv^2)}{dr} \right)^2. \quad (8)$$

$$2r \frac{d^2 v}{dr^2} + 5 \frac{dv}{dr} + \frac{4r}{v} \left( \frac{dv}{dr} \right)^2 + \frac{A}{4\pi G} \left( \frac{v^2}{r^2} + \frac{4v}{r} \frac{dv}{dr} + 4 \left( \frac{dv}{dr} \right)^2 \right) = 0. \quad (9)$$



## ***Solution of the equations***

• This nonlinear differential equation of the second order has a singularity at the origin, which is a generally recognized feature and difficulty in the hydrodynamic interpretation of rotational curves. However, by lowering the order of the equation, we have circumvented this difficulty. If we substitute  $v = C \exp \int q dr$ , into equation (9) where  $q$  is an unknown function,  $C$  is a constant. Then from (9) we obtain an equation for  $q$ , which we consider a smooth function (  $\frac{dq}{dr} = \frac{d^2q}{dr^2} = 0$  ). In this approximation

$$q \frac{AC}{4\pi G} = - \frac{2q^2}{\left( \frac{1}{r^2} + \frac{4}{r}q + 4q^2 \right)} + \frac{(6rq^2 + 5q) \left( -\frac{2}{r^3} + \frac{4}{r^2}q \right)}{\left( \frac{1}{r^2} + \frac{4}{r}q + 4q^2 \right)^2} \quad (10)$$

## Problem solution


Keeping only the leading terms close  $r \rightarrow 0$ , we find that approximately, taking into account the two roots of equation (10) and the choice of constants, when the velocity vanishes at  $r \rightarrow 0$

$$v \approx C \sin \left( \frac{\sqrt{2 + \frac{4\pi G}{AC} 60r}}{12} - \frac{\sqrt{2}}{12} \right) \quad (11)$$

which depends on the choice of constant  $C$ . We find the constant  $C$  by comparing the solution (11) for the transverse velocity with the experimental rotation curves of our Galaxy [11]. In our case, it turns out that  $C \approx 200 \text{ km/s}$ . With this choice of constant  $C$ , we manage to reproduce the increase in velocity  $v$  with distance  $r$  from the center of the Galaxy due to dark matter. In this case, as the constant  $\langle \sigma v \rangle$  we use the value  $10^{-20} \text{ cm}^3/\text{s}$  [12] and  $A = 3 \cdot 10^6 \text{ cm}^3/\text{s/g}$ , the constant  $G = 6.67 \cdot 10^{-11} \text{ m}^3/\text{kg/s}^2$ . With a mass of dark matter particles of the order of 10-100 MeV, or more precisely 17 MeV, expression (11) agrees with the available experimental data [11].



Figure shows the dependence of the velocity  $v$  on the distance  $r$  according to formula (11) (red line), the contribution to the transverse velocity due to attraction to the center of the Galaxy without taking into account dark matter (blue dashed line) (contribution of the bungee and disk), the resulting curve (blue line) taking into account the contribution of dark matter according to (11), red dots are experimental data from [11].



## DISCUSSION

- As a result of comparison with the experimental slope of the rotation curve, it turned out that this corresponds to a dark matter particle mass of about 10-100 MeV. This is close in mass to the new particles X17 and X38 with masses of 17 MeV and 38 MeV, respectively. The interpretation [6,7] of the detection of which at accelerators and their discovery by the Hungarian group ATOMKI [6,7] in Be-8 decays were discussed by us earlier [13,14]. The fact that the mass of dark matter particles turns out to be about 17 MeV can also be seen from equation (7).
- Indeed, knowing the average density of matter in our Galaxy  $\rho \approx 1.8 \cdot 10^{-24} \text{ kg/cm}^3$ , for  $r \approx 3 \cdot 10 \text{ kpc}$  we obtain
- $$v \approx Apr \approx 10^{-20} / (17 \cdot 1.6 \cdot 10^{-27} \cdot 10^{-3}) 1.8 \cdot 10^{-24} \cdot 3 \cdot 10 \cdot 10^{16} \text{ km/s} \approx 200 \text{ km/s}$$
- A similar assessment is given in [15] The matter remains to be seen in subsequent experiments.



## *DISCUSSION of New particles*

- The X17 boson predicted in Refs [6, 7] may manifest itself in the spectrum of soft photons. This was reported in Wong's papers [8, 9] as well as in our papers [1-3]. Wong [16] proposed an interpretation of this boson as a result of combining QCD and QED. In this case, the union is performed for two-dimensional QCD<sub>2</sub> QED<sub>2</sub> in the tube model
- In [1], we proposed a modified tube model. In this case, as well as Wong [16], we consider both the hadronic tube at the coupling constant
- $\alpha = \alpha_s$  and the electromagnetic tube at  $\alpha = \alpha_e = 1/137$ . The tube radii are determined from the minimum energy per unit length, and the strong interaction constant is  $\alpha_s \approx 0.5$ . According to the model of a tube stretched between two quarks [1], one can find the masses of the resulting hadrons, and in the case of an electromagnetic tube, the mass of the X17 boson. In this case, according to [18, 19], the energy density of the tube consists of two terms:

$$\rho = A + G \quad (12)$$

- The first term  $A$  is determined by the field strength  $E$ . If the field were distributed uniformly, then the energy per unit length of the tube of radius  $r$  would be distributed into the hemisphere, equal to

$$A = E^2 \frac{2\pi}{4\pi} \pi r^2 = \frac{1}{2} E^2 \pi r^2 \quad (13)$$

But, since in the direction of the length of the tube it is necessary to sum only the solid angles multiplied by  $\cos \theta$ , the projections of the unit vector onto the direction of the tube, then the solid angle

$$\Omega = \int_0^{\pi/2} \cos \theta \sin \theta d\theta \int_0^{2\pi} d\varphi = \pi \quad (14)$$

but not  $2\pi$ . That's why

$$A = E^2 \frac{\Omega}{4\pi} \pi r^2 = \frac{1}{4} E^2 \pi r^2, \quad (15)$$

where the intensity vector

$$E \text{ flux is } \Phi = \int_0^{\pi/2} E \cos \theta d\theta \int_0^r 2\pi r dr = E \pi r^2, \text{ and intensity}$$

$$E = \frac{\Phi}{\pi r^2} = \frac{g}{\pi r^2} \text{ and the coupling constant } \alpha = \frac{g^2}{4\pi}, \text{ } r \text{ is the radius of the tube.}$$

$$\text{Hence } A = \frac{\alpha}{r^2} \quad (16)$$

The second term is expressed in terms of the bag constant  $B = 0.17 \text{ GeV/fm}^3$  [12]:

$$G = B \pi r^2 \frac{\alpha}{\alpha_s} \quad (17)$$

where we introduced the ratio of the constant  $\alpha$  to the strong interaction constant  $\alpha_s$ . Here  $\text{fm} = 10^{-15} \text{ m}$



- For an oscillating rectilinear string-tube, we obtain [10] the mass :

$$M^2 = 2\pi\rho n \quad (18)$$

- where  $n$  is the quantum number. For the hadron tube at  $n=1$ , we obtain
- $M \approx 152 \text{ MeV}$  for the  $\pi^0$ -meson. But if the bag constant is chosen equal to  $0.13 \text{ GeV/fm}^3$ , then for the  $\pi^0$ -meson we obtain a  $M \approx 140 \text{ MeV}$  value that is closer to the experimental one. For an electromagnetic tube with the same tube radius, we obtain the mass of the neutral **X17 boson 17 MeV**. According to the formula  $M^2 = 2\pi\rho m$ , resonances can also be obtained, where  $m$  is a multiple folded string with rotation. So you can get the  $\rho$  mass - meson, and for the electromagnetic string we get the mass of the **X38 boson 34 MeV** at  $m=4$  obtained in Wong's work in a different way.
- Note that these results were obtained in our approach using formulas different from Wong's. In his work [16], Wong proposes to interpret the **X17 boson as a dark matter particle, since it is neutral, not a baryon, and can be a composite particle of large-mass astrophysical objects.**

# Ultrahigh energy cosmic ray

In this paper, it is proposed to check the manifestation of these new particles in the spectra of cosmic rays of ultrahigh energies of the order of  $10^{19}$  eV. Indeed, bursts in the spectrum of cosmic rays in the energy region of  $10^{11}$  GeV were obtained in [21, 22], which are interpreted in [23] as a manifestation of a dark photon. This dark photon has a mass different from zero, but very small, less than  $10^{-6}$  eV.

We propose an interpretation of these bursts as a contribution from the decay of new X17 and X38 particles in the framework of our approach [1-3]. Indeed, the contribution from the decay of the X boson into photons can be approximately represented as

$$\frac{dN}{dE} \propto p \exp \left( - \frac{\sqrt{m^2 + (p - p_0)^2} - m}{T} G \right), \quad (19)$$

where  $m$  is the mass of the X-boson,  $p$  is the momentum,  $T$  is the temperature,  $G$  is the effective Lorentz factor, taking into account the motion of X bosons in cosmic rays,  $p_0$  is the position of the burst.

In order to determine the temperature  $T$  and the factor  $G$ , we use an approximation of the cosmic radiation spectrum before the burst

$$\frac{dN}{dE} \propto \exp \left( - \frac{p}{T} G \right) \quad (20)$$

which is known and approximately proportional to  $E^{-3}$ . From here we find  $G$ , assuming that the temperature  $T \approx 14$  MeV is about 10 times less than the temperature of hadrons 140 MeV, since we are interested in electromagnetic interactions. The same is true and of our X-bosons. Included in formula (8) is found from the definition of the peak maximum.

Approximately  $p - p_0 \approx T / G$  relative to the tail of the spectrum (9) for small  $m$ . As a result, we compared this with experimental data. We compared the experimental points from [13, 14] and the data from [15] for an ultralight dark photon



$$E^3 dN/dE, \text{ GeV}^2/(\text{m}^2 \text{ s sr})$$




Figure. Spectrum of ultrahigh-energy cosmic rays. Experimental points from [21, 22]. The solid curve 1 is the spectrum from a dark photon from [23]. Solid curve 2 is our spectrum of photons from the X17 boson decay in accordance with formula (19), dashed curve without taking into account the boson decay according to formula (20), Solid curve 3 corresponds to a mass of 1000 MeV, curve 4 corresponds to a mass of 10 GeV, dashed curve 5 corresponds to mass 100 GeV

Curve 1 from

P, Tantirangsri, D. Samart, C. Pongkitivanichkul. ArXiv 2301.11122 v1[hep-ph].

Points from

M. Takeda, N. Hayashida, K. Honda et al. Phys.Rev. Lett. 81, 1163 (1998).

D.J. Bird, S.C. Corbató, H.Y.Dai et al. Phys.Rev. Lett. 71, 3401 (1993).

## CONCLUSION

- Thus, we are convinced that one more confirmation of the existence of X17 and X38 bosons with masses of 17 MeV and 38 MeV has been obtained in the light of experimental data from cosmic rays for ultrahigh energies that are not achievable with modern accelerators. Heavy particles of large mass also do not contradict these experimental data.
- As a result of comparison with the experimental slope of the rotation curve, it turned out that this corresponds to a dark matter particle mass of about 10-100 MeV. This is close in mass to the new particles X17 and X38 with masses of 17 MeV and 38 MeV, respectively



# References

1. A.T. D'yachenko, Phys. Atom. Nucl. 83, 1597( 2020).
2. A.T. D'yachenko, Phys. Atom. Nucl. 85, 1028(2022)
3. A.T. D'yachenko, A.A. Verisokina, M.A. Verisokina, Acta Phys. Polonica. B Proc. Suppl. 14, 761(2021).
4. A. Belogianni *et al.*, Phys. Lett. B548, 129 (2002).
5. K. Abraamyan, C. Austin, M. Baznat, *et al.*, EPJ Web of Conf. 204, 08004 (2019); arXiv: 1208.3829
6. A.J. Krasznahorkay, M. Csatós, L. Csige, *et al.*, Phys. Rev. Lett. 116, 042501 (2016); arXiv: 1504.01527
7. A.J. Krasznahorkay, M. Csatós, L. Csige, *et al.*, Phys. Rev. C 104, 044003 (2021); e-Print 2104.10075 [nucl-ex].
8. А.Г. Дорошкевич, В.И. Лукаш, Е.В. Михеева, УФН, 182, 5 (2012).
9. В.С. Березинский, В.И. Докучаев, Ю.Н. Ерошенко, УФН, 184, 3 (2014).
10. J.G. Mamprim<sup>1</sup>, G. Gambini, L. B. Arbeleche<sup>1</sup>, M. Olegario, V. Souza<sup>1</sup>;  
<https://arxiv.org/abs/2502.11958>
11. F. Combes, J. Lequeux, The Milky Way: Structure, Dynamics, Formation and Evolution.  
— Les Ulis: [EDP Sciences](#), 2016. — 195 p. — [ISBN 978-2-7598-1915-7](#).
12. R. Brant, Z. Andrew Phys. Rev. D 79 083525 (2009)
13. А.Т. Дьяченко, ЯФ, 86, 259 (2023)
14. М.В. Архипов, А.Т. Дьяченко, Письма в ЖТФ, 50 (23), 11(2024).
15. B. B. Kamenetskaia, J.-C. Park, M. Reichard, and G. Tomar, Investigating sub-MeV dark matter annihilation to neutrinos using direct detection experiments. arXiv:2505.07403v1 [hep-ph] 12 May 2025
16. C.Y. Wong, JHEP, 08165 (2020); arXiv: 2001.04864v1 [nucl-th]
17. C.Y. Wong, Phys. Rev. C 81, 064903 (.2010).
18. V.A. Abramovsky, E.V. Gedalin, E.G. Gurvich, and O.V. Kancheli. *Inelastic interactions at high energies and the chromodynamics*. Tbilisi: Metsniereba, 1986.
19. B.M. Barbashov and V.V. Nesterenko. *Relativistic string model in hadron physics*. - M. Energoatomizdat, 1987.
20. V.M. Emelyanov, S.L. Timoshenko, and M.N. Strikhanov. *Introduction to relativistic nuclear physics*. - M.: Fizmatlit, 2004
21. M. Takeda, N. Hayashida, K. Honda *et al.* Extension of the cosmic-ray energy spectrum beyond the predicted Greisen-Zatsepin-Kuz'min cutoff // Phys.Rev. Lett. 81, 1163 (1998).
22. D.J. Bird, S.C. Corbató, H.Y. Dai *et al.* Evidence for correlated changes in the spectrum and composition of cosmic rays at extremely high energies. // Phys.Rev. Lett. 71, 3401 (1993)
23. P. Tantirangsri, D. Samart, C. Pongkitivanichkul. Dark photon bremsstrahlung and ultra-high –energy cosmic ray. ArXiv 2301.11122 v1[hep-ph].



THANK YOU !



22 Lomcon MSU 26 August 2025